# Stuarts Draft, VA

# **Project Summary**

As part of Virginia's State-Based Anemometer Loan Program, an anemometer was placed in Stuarts Draft, Virginia to assess the area's wind energy potential. The data presented reflects wind speed and direction data recorded from March 1, 2005 through February 28, 2006. Annual average wind speed during the monitoring period (66 ft. [20 m] above ground level) was recorded to be 7.42 mph (3.31 m/s). Wind power density is calculated to be 65.17 W/m<sup>2</sup>.

# **Project Location**

The monitoring equipment was installed on private land in Stuarts Draft, Virginia at an elevation of 1,500 feet. The site is located at N 38° 5.16', W 79° 45.84'.

# **Monitoring Equipment**


The 20-meter NRG-NOW Systems – Wind Explorer kit includes one tower, one Wind Explorer data logger with shelter box, one #40 Maximum anemometer, one #200P wind direction vane, sensor cabling, one lightning rod with copper ground, and two data plugs.

#### Results

Average annual wind speed	7.42 mph (3.31 m/s)
Average annual wind power density	65.17 W/m <sup>2</sup>
Month with greatest wind resource	February
Average wind speed during month with greatest resource	10.8 mph
Month with least wind resource	August
Average wind speed during month with least resource	4.4 mph


#### **Monthly Variation**

Figure 1 below shows the average monthly wind speed for each month throughout the recorded year. Typical sites see high values in the winter months; this site however, does not follow such a trend, which may be applicable to localized shading from surrounding trees.



#### **Diurnal Variation**

Figure 2 shows how wind speeds vary on an hourly basis. Stronger winds are more prevalent during late morning and afternoon hours. Slower winds more often exist after midnight and during sunrise.



#### **Monthly Diurnal Variations**

Figure 3 shows the diurnal (hourly) variations of the wind speed over each month. The trends of each month fairly closely follow one another for this site.




Figure E.3.3. Average hourly wind speeds for each month.

#### Wind Speed Histogram

Figure 4 shows the occurrence in number of hours that each wind speed occurs. Typically this falls under what is called the Weibull distribution, with a high peak early, and a tailing off at higher wind speeds.

Data (Mar-05 to Feb-06)




Figure 4. Wind speed frequency over the year by # of hours.

#### Wind Speed and Direction Trace

Figure 5 shows the wind speed and wind direction reading for every ten-minute average over the course of the data. The wind speed data is the blue line, and the directional data are the red lines.




Figure 5. Wind speed and direction readings throughout the year.

#### Wind Daily Averages

Figure 6 shows the average wind speed and direction by day of the month. Notice there is no particular trend here.




Figure 6. Wind speed and direction averages by day of the month.

#### Wind Speed Occurrence by Direction

Figure 7 displays the average wind speed for each direction the wind comes from. It also shows the percent of time in that direction bin. For this site the higher wind speeds came from the southwest, as well as the majority of the winds.




Figure 7. Average wind speed for each direction bin and % time in that bin.

#### **Frequency Wind Rose**

Figure 8 displays a frequency wind rose. It can be seen that the majority of the wind come from the southwest.




Figure 8. Wind rose displaying wind direction frequency.

# **Energy Rose**

Figure 9 is an energy wind rose, showing the number of  $kWh/m^2$  available from 16 compass-point directions. The majority of energy comes from the southwest. This is due to the high wind speeds and high frequency of occurrence of these winds.




Figure 9. Energy wind rose with units in  $kWh/m^2$ .